



**University of  
Zurich<sup>UZH</sup>**

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2011

---

## **A flux-sensing mechanism could regulate the switch between respiration and fermentation**

Huberts, Daphne H E W ; Niebel, Bastian ; Heinemann, Matthias

**Abstract:** The yeast *Saccharomyces cerevisiae* can show different metabolic phenotypes (e.g. fermentation and respiration). Based on data from the literature, we argue that the substrate uptake rate is the core variable in the system that controls the global metabolic phenotype. Consequently the metabolic phenotype that the cell expresses is not dependent on the type of the sugar or its concentration, but only on the rate at which the sugar enters the cell. As this requires the cells to 'measure' metabolic flux, we discuss the existing clues toward a flux-sensing mechanism in this organism and also outline several aspects of the involved flux-dependent regulation system. It becomes clear that the sensing and regulation system that divides the taken up carbon flux into the respiratory or fermentative pathways is complex with many molecular components interacting on multiple levels. To obtain a true understanding about how the global metabolic phenotype of *S. cerevisiae* is controlled by the glucose uptake rate, different tools and approaches from systems biology will be required.

DOI: <https://doi.org/10.1111/j.1567-1364.2011.00767.x>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-79182>

Journal Article

Originally published at:

Huberts, Daphne H E W; Niebel, Bastian; Heinemann, Matthias (2011). A flux-sensing mechanism could regulate the switch between respiration and fermentation. *FEMS Yeast Research*, 12(2):118-128.

DOI: <https://doi.org/10.1111/j.1567-1364.2011.00767.x>

# Rate of sugar uptake determines the global metabolic phenotype of yeast

Daphne H. E. W. Huberts, Bastian Niebel, Matthias Heinemann

Molecular Systems Biology, Groningen Biomolecular Sciences and Biotechnology Institute,  
University of Groningen, Nijenborgh 4, 9747 AG Groningen, The Netherlands

Corresponding author: m.heinemann@rug.nl (phone +31 50 363 8146)

## Abstract

The yeast *S. cerevisiae* can show different metabolic phenotypes (e.g. fermentation and respiration). Based on data from the literature, we argue that the substrate uptake rate is the core variable in the system that controls the global metabolic phenotype. Consequently the metabolic phenotype that the cell expresses is not dependent on the type of the sugar or its concentration, but only on the rate at which the sugar enters the cell. As this requires the cells to ‘measure’ metabolic flux, we discuss the existing hints towards flux-sensing mechanism in this organism and also outline several aspects of the respectively involved flux-dependent regulation system. It becomes clear that the sensing and regulation system that divides the taken up carbon flux into the respiratory or fermentative pathways is complex with many molecular components interacting on multiple levels. To obtain a true understanding about how the global metabolic phenotype of *S. cerevisiae* is being realized, different tools and approaches from systems biology will be required.

## Introduction

Microorganisms are constantly facing changing environments, for example, in terms of their nutrient availability. In order to be informed about the characteristics of the environment, sensory systems are required. Classically, these are either transmembrane receptors (Holsbeeks *et al.*, 2004; Rubio-Teixeira *et al.*, 2010) or intracellular receptors (such as transcription factors) whose activity is modulated by

their specific ligands. While for certain substrate molecules we know such sensory systems (in yeast e.g. the transmembrane receptors Snf3p and Rgt2p (Johnston & Kim, 2005), or the intracellular receptor Gal3p (Sellick & Reece, 2005; Campbell *et al.*, 2008), for many other substrates we have not yet identified specific sensors. The question is whether we simply do not know them yet or whether cells recognize these metabolites in a different way.

An alternative way to sense the presence of a certain carbon source would be by measuring the metabolic flux that derives from its degradation. In this way cells could perceive what substrate is being imported and at which rate. However, all sensing mechanisms in biology are based on concentration measurements, so cells would need to translate a rate (i.e. flux) into concentrations of certain flux-signaling molecules. Actually, it has been shown for *E. coli* in a synthetic, engineered system that it is possible for cells to ‘measure metabolic fluxes’ and use them for regulation (Fung *et al.*, 2005). In addition, a recent computational study that modeled *E. coli* central metabolism suggested that intracellular metabolic fluxes is indeed used to indirectly perceive the presence of a particular carbon source and the rate with which it enters the cell (Kotte *et al.*, 2010). Also for yeast, it was occasionally proposed that it can measure metabolic flux (Ye *et al.*, 1999; Bisson & Kunathigan, 2003; Agrimi *et al.*, 2011), but it was similarly often explicitly stated that this would not be possible (Gamo *et al.*, 1994; Rodriguez & Flores, 2000; Youk & van Oudenaarden, 2009).

In this perspective article, we explore flux-sensing and flux-dependent regulation in yeast. Specifically, we will look (i) at what kind of evidence or indication we have for flux-dependent regulation, (ii) at general conceptual issues with sensing a rate and at potential flux-sensing mechanisms in yeast, and (iii) at some properties of the regulatory machinery that controls metabolism in a flux-dependent manner. As flux-sensing and flux-dependent regulation will not be accomplished via a single molecule, but rather will be realized by a whole system of molecules interacting in a very specific and likely complicated manner, systems biology approaches and modeling will definitely be required to generate a comprehensive understanding about this system. Thus, we conclude with a brief discussion about how systems biology could be instrumental towards this goal.

## I. Evidence for metabolic flux-dependent regulation

An indication for flux-dependent regulation would be a correlation between the magnitude of a certain metabolic flux and a particular phenotype. Within limits, chemostat cultures allow modulation of metabolic fluxes, while at the same time keeping environmental conditions almost constant (Weusthuis *et al.*, 1994; Bull, 2010). A flux/phenotype correlation can be seen, for example, between the glucose uptake rates and the extent of usage of the fermentative pathway, with specifically particularly the ethanol excretion rate increasing with the glucose uptake rate. This is evidenced data from aerobic glucose-limited chemostat cultures of *S. cerevisiae* (cf. inset of Fig. 1A with data from Cortassaa & Aon, 1998; van Hoek *et al.*, 1998; Diderich *et al.*, 1999; Daran-Lapujade *et al.*, 2004; Frick & Wittmann, 2005; Daran-Lapujade *et al.*, 2007; Jouhten *et al.*, 2008; Basso *et al.*, 2010, cf. also Supplementary information).

In contrast to limiting the glucose uptake rate through environmental conditions, manipulating the maximal glucose uptake rate by genetic means is an alternative approach to assess the importance of the glucose uptake rate in dictating the metabolic mode. Elbing *et al.* (2004) constructed a set of hexose transporter mutant strains that only differ in their maximal glucose uptake rate (Elbing *et al.*, 2004). Similar to results obtained in glucose-limited chemostat cultures, it was found that the different transporter strains metabolize glucose to various extents by respiration and fermentation dependent on the rate of glucose uptake in an identical (i.e. high glucose) environment (Elbing *et al.*, 2004). This data perfectly aligns with the above mentioned chemostat data (cf. inset of Figure 1A and Supplementary information). For example, the phenotype of the strain with the lowest glucose uptake capability (TM6\*) when grown at high glucose is the same as the one of the respective wildtype in the chemostat culture at a low dilution rate when the glucose influx is restricted by the environmental conditions. As in these experiments and in similar studies generalizing the findings from Elbing *et al.* to other *S. cerevisiae* strains (Reifenberger *et al.*, 1995; Ye *et al.*, 1999; Henricsson *et al.*, 2005), the environment was identical and the strains used differ only in their glucose transport capability, the glucose uptake rate seems to be the control factor for the metabolic mode and not environmental conditions such as the extracellular glucose concentration, which is inherently different in batch and

glucose-limited chemostat cultures. Indeed, also glucose sensing mutants switch from respiration to fermentation at similar glucose uptake rates as their corresponding wildtype strain (cf. inset of Figure 1A with data from (Cortassa & Aon, 1998). Also data from various different *S. cerevisiae* strains grown in glucose batch cultures fit nicely to the so far presented data (cf. inset of Figure 1A with data from Blom *et al.*, 2000; Peter Smits *et al.*, 2000; Elbing *et al.*, 2004; Otterstedt *et al.*, 2004; Blank *et al.*, 2005; Cordier *et al.*, 2007; Velagapudi *et al.*, 2007; Heyland *et al.*, 2009; Kummel *et al.*, 2010; Christen & Sauer, 2011; Costenoble *et al.*, 2011; Raab *et al.*, 2011).

As the correlation between glucose uptake rate and ethanol secretion rate seems to be independent of the culturing methods (e.g. chemostat, batch cultures), the glucose concentration, and the *S. cerevisiae* strains used, we set out to challenge this correlation further. In absence of oxygen yeasts are forced to ferment, and this condition may uncouple glucose uptake rate from ethanol secretion rates. However, even though under anaerobic conditions the ethanol production rate is generally higher as under aerobic conditions (Verduyn *et al.*, 1990; Peter Smits *et al.*, 2000; Aguilera *et al.*, 2005; Jouhten *et al.*, 2008; Wiebe *et al.*, 2008), the anaerobic data fit well to the correlation presented in the inset of Figure 1A (cf. Figure 1A). Along the same line, data from yeast grown under various environmental conditions, such as different pH or salinity of the medium (Heyland *et al.*, 2009), temperature (Tai *et al.*, 2007; Postmus *et al.*, 2008; Heyland *et al.*, 2009), addition of weak organic acids (Larsson *et al.*, 1997; Abbott *et al.*, 2007; Daran-Lapujade *et al.*, 2007) and nitrogen limited chemostats (Larsson *et al.*, 1997; Meijer *et al.*, 1998; Diderich *et al.*, 1999), also align well with the already plotted data (cf. Figure 1A). Even when data from a wide range of *S. cerevisiae* mutant strains is added, the correlation between glucose uptake rate and ethanol secretion rate still remains intact. Examples of such mutant strains include strains with overexpressed enzymes of lower glycolysis (Peter Smits *et al.*, 2000), uracil auxotrophies (Basso *et al.*, 2010), single deletions or overexpressions of various proteins involved in glucose regulation (Blom *et al.*, 2000; Raab *et al.*, 2011) and carbon metabolism (Blank *et al.*, 2005; Cordier *et al.*, 2007; Velagapudi *et al.*, 2007) (cf. Figure 1A).

Due to this robustness of the correlation between the glucose uptake and ethanol secretion rates, we next asked whether this correlation would also be maintained in different yeast species. Yeast species

can roughly be subdivided in Crabtree positive and Crabtree negative yeast (De Deken, 1966). Although the term “Crabtree effect” has been used with different meanings (Barford & Hall, 1979; Van Urk *et al.*, 1990), ‘Crabtree positive’ means here that a certain yeast species can produce ethanol under aerobic conditions, which ‘Crabtree negative’ cannot (De Deken, 1966; Verduyn *et al.*, 1992). It is sometimes argued that Crabtree negative yeasts have a higher respiratory activity than Crabtree positive yeasts and, as a result, Crabtree negative yeast would not have to resort to ethanol formation under high glucose conditions (Verduyn *et al.*, 1991). However, when data from various Crabtree positive and negative yeast species (gathered from Christen & Sauer, 2011; Rozpedowska *et al.*, 2011) is plotted in Figure 1A, it turns out that the measured glucose uptake rates and ethanol secretion rates fall perfectly on the already established correlation, indicating that Crabtree negative yeast species have simply a lower glucose import rate and thus show no fermentative activity; in line of what was also suggested earlier (Does & Bisson, 1989; van Urk *et al.*, 1989; Boles & Hollenberg, 1997; Rozpedowska *et al.*, 2011). This idea is further supported by the observation that Crabtree negative yeast species (*Candida utilis*, *Hansenula polymorpha* and *Kluyveromyces marxianus*) can ferment under aerobic conditions by stimulating glucose uptake using a weak organic acid (Verduyn *et al.*, 1992) (data not shown).

Next we asked whether the correlation also holds when other sugars are taken up instead of glucose. Generally, *S. cerevisiae* ferments glucose, but other sugars, such as sucrose, mannose and galactose, are respired or fermented to different extents (Fendt & Sauer, 2010). Based on our established correlation between glucose uptake and ethanol secretion rate, we asked whether the capability to grow fermentatively on certain sugars is determined only by the respective import rate that yeast can realize with a certain sugar. Indeed ethanol production rates obtained from *S. cerevisiae* grown on galactose (Sierkstra *et al.*, 1993; Diderich *et al.*, 1999; Ostergaard *et al.*, 2000; Velagapudi *et al.*, 2007; Costenoble *et al.*, 2011) compare well with those measured for glucose at the same sugar uptake rate (see Figure 1A). This is also true for other sugars, such as fructose (Diderich *et al.*, 1999), maltose (Weusthuis *et al.*, 1994; Wisselink *et al.*, 2007; de Kok *et al.*, 2011), arabinose (Wisselink *et al.*, 2007) and sucrose (Basso *et al.*, 2010), when normalized to c-mols (see Figure 1A and Supplementary

information). Beyond, a relationship between import rate and the ability to ferment a particular sugar was also seen for maltotriose (Zastrow *et al.*, 2001; Dietvorst *et al.*, 2005) and cellobiose (Ha *et al.*, 2011). Identical to glucose, a strain with an engineered higher galactose uptake rate increased its respiro-fermentative metabolism with the ethanol production rate increasing linearly with glycolytic flux (Ostergaard *et al.*, 2000). Similarly, *Kluyveromyces lactis* switched to aerobic fermentation of galactose after introduction of the GAL2 gene (Goffrini *et al.*, 2002). In contrast, a yeast strain lacking invertase activity and with only a limited capacity to transport sucrose into the cell, showed a significantly reduced ethanol production rate (Badotti *et al.*, 2008). Overall, these observations indicate that the distribution between the respiratory and fermentative pathways is dependent not on the type of substrate being consumed but by the rate at which that substrate is imported.

The above mentioned observation that the ethanol excretion rate generally correlates with sugar uptake rate points towards a flux-dependent regulation. It could, however, also be conceivable that the growth rate determines the physiology, as, for example, was suggested by van Hoek *et al.*, 1998. To test this hypothesis, we plotted the ethanol excretion rates against the respective growth rates. As we here find absolutely no correlation (cf. Figure 1B), growth rate is very unlikely to be the determining control factor.

Another potential issue could be that the identified correlation could simply also be caused by a physical rate limitation somewhere in metabolism. Indeed it has been argued that the ethanol production is due to an overflow mechanism (Sonnleitner & Kappeli, 1986; van Hoek *et al.*, 1998; Zhuang *et al.*, 2011) meaning that there might a limitation in the TCA cycle or respiratory chain that causes the excretion of ethanol with increasing sugar uptake. If this would be the case, ethanol excretion would not be the result of an active, flux-dependent regulation. However, there are a number of indications that speak against the overflow hypothesis: (i) If the ethanol excretion would be the result of a capacity limitation in the respiratory chain, i.e. respiration reaches a maximum level at some point and no further NADH generated in the TCA cycle could be respired, then we would expect the oxygen uptake rate to stay at a constant (high) level with further increasing sugar uptake rates. As usually a decrease in the O<sub>2</sub> uptake rates is reported with increasing glucose uptake rates (Beck & von

Meyenburg, 1968; Verduyn *et al.*, 1992; van Hoek *et al.*, 1998; Canelas *et al.*, 2011), it seems that at least respiration could not be the limiting factor. Further, the highest possible O<sub>2</sub> uptake rate can also not be reached, which becomes obvious if we compare the O<sub>2</sub> uptake rate when yeast is grown on ethanol with the highest O<sub>2</sub> uptake rate reported on when yeast grows on glucose (7.2 mmol O<sub>2</sub>/gDW/h in aerobic glucose chemostat culture at growth rates between 0.25 1/h and 0.33 1/h (Canelas *et al.*, 2011) compared to 11.8-13 mmol O<sub>2</sub>/gDW/h for unlimited growth on ethanol (as estimated on the basis of measured ethanol uptake rates (Costenoble *et al.*, 2011) and flux variability analysis (Mahadevan & Schilling, 2003)). (ii) It was reported that an alcohol dehydrogenase (*ADHI*) deletion mutant in *S. cerevisiae* is very sick on high glucose conditions - a condition requiring fermentation. In contrast, the *ADHI* deletion has no phenotype on galactose - a condition, on which *S. cerevisiae* only respire. In turn, deletion of the TCA cycle isoenzyme succinate dehydrogenase (*SDHI*) has no phenotype on glucose, but a lethal one on galactose (Ewald, Matt, Zamboni, 2011, unpublished results). Similarly, blocking respiration on substrates that are typically respired leads to no growth; a phenotype that can be rescued by overexpressing of the sugar transporters (Goffrini *et al.*, 2002; Fukuhara, 2003). Obviously, the sugar uptake rate and the metabolic mode need to fit together. In our opinion, altogether these findings point more to a flux-dependent regulation than to a simple physical overflow mechanism. If the latter would be true, then situations where the TCA cycle or the respiration is blocked or reduced could simply be rescued by fermenting the galactose.

As a result, we propose that there is a universal regulatory system in place that (i) measures flux and then (ii) determines the ratio in which glucose is utilized by respiration or fermentation according to an evolutionary conserved program (see Figure 2A).

## **II. How could metabolic flux be sensed?**

### **a. Theoretical considerations**

We have seen that the global metabolic phenotype assumed by yeast is dependent on the substrate uptake rate, i.e. that there is flux-dependent regulation. This implies that a flux signal needs to be



sensed within the cell. Before we discuss how a cell could actually achieve this, we first introduce – from a theoretical viewpoint – two concepts of flux-sensing.

A flux through a metabolic reaction is nothing else than a reaction rate. Here a rate  $r$  is defined as the ratio between an infinitesimally small change of a state quantity  $x$ ,  $dx$ , and an infinitesimally small change in time  $t$ ,  $dt$ , according to

$$r = \frac{dx}{dt} .$$

In other words, a rate is a derivative of a state variable. The first way to sense a rate  $r$ , is by measuring the difference of state  $x$  between two time points  $t$ ,

$$r \approx \frac{\Delta x}{\Delta t} .$$

If the cell would follow this concept, it would need to have the capability to memorize two states, to measure time and to do mathematical operations in terms of subtractions and a division. It is obvious that cells cannot do this. However, there is a second possibility of how cells could get informed about a rate  $r$ . A rate  $r$  can be estimated from a state  $x$  (i.e. a metabolite concentration), if the functional dependencies of the system  $f$  are known, according to

$$r = f(x) .$$

This concept requires that the cell ‘knows’ the system that generates the state  $x$ , i.e. it has a model of the functional dependencies of the system,  $f$ . For cells to use this concept, it is important that a simple (ideally linear) dependency exists between  $r$  and  $x$  and that no other state variables influence this relationship as the controller (i.e. the regulatory system) that receives the flux-signal  $x$  needs to ‘work’ with this signal. Thus, cells that exploit flux-dependent regulation need to have a system that translates a flux ( $r$ ) into a concentration of a biomolecule ( $x$ ). This, for example, flux-signaling metabolite  $x$  then would induce a flux-dependent regulation (see Figure 2B).

## **b. Hints towards flux-sensing mechanisms in yeast**

Where in the metabolic network could the flux be sensed? The fact that the ethanol excretion rate not only correlates with the glucose uptake flux, but with all sugars' uptake rates (cf. Figure 1A) suggests that the flux sensor needs to be at a point in metabolism that is equally affected by all such sugars. The here mentioned sugars converge at either glucose 6-phosphate (G6P) (glucose, maltose, galactose), at fructose 6-phosphate (F6P) (fructose) or at both (sucrose). Thus, the flux sensing mechanism resides likely below F6P. On the basis of this reasoning, specific transporters or hexokinase 2 (Hxk2) can be excluded, although the latter is frequently suggested to be involved in flux-sensing (Bisson & Kunathigan, 2003).

If we argue here that eventually the flux is measured in glycolysis somewhere below F6P, and we had earlier only established a correlation between the sugar uptake rate and the global metabolic phenotype, then we need to check a correlation also exists between the sugar uptake rate and the glycolytic flux. A correlation is not necessarily expected, because glucose (or G6P) is also shuffled into the pentose phosphate pathway and into storage metabolism. Nevertheless, data from (partly <sup>13</sup>C based) metabolic flux analyses demonstrate that the glycolytic flux between G6P and F6P linearly correlates with the sugar uptake rate (see Figure 3 with data from Nissen *et al.*, 1997; Gombert *et al.*, 2001; Blank *et al.*, 2005; Frick & Wittmann, 2005; Jouhten *et al.*, 2008; Fendt & Sauer, 2010; Christen & Sauer, 2011) for glucose and galactose independent of the yeast species and the culture conditions employed and that this correlation is also robust against many environmental and genetic perturbations.

One way to establish a relationship of the kind  $r=f(x)$  would be with  $x$  being the concentration of a metabolite that would correlate (ideally) linearly with the flux  $r$ . As metabolite concentrations were found to be highly specific for the limiting nutrient (Boer *et al.*, 2010), a signaling role for metabolites would not be so farfetched. Remarkably, the concentration of fructose-1,6 biphosphate (FBP) seems to correlate with the sugar uptake rate when *S. cerevisiae* data from glucose batch (Fendt *et al.*, 2010) and glucose-limited chemostat cultures with different dilution rates (Canelas *et al.*, 2011) and different cultivation temperatures (Postmus *et al.*, 2008) are plotted (see Figure 4A). This correlation even holds for data from other Crabtree positive and negative yeast species (Christen & Sauer, 2011) (see Figure

4A). Thus, FBP could be a flux-signaling metabolite, like it was also suggested to be in *E. coli* (Kotte *et al.*, 2010). Interestingly, when the glucose concentration is suddenly increased in a glucose-limited chemostat culture or glucose is added to an ethanol-limited chemostat culture, the glucose influx rate increases with a concomitant increase in the concentration of FBP and onset of ethanol excretion. This indicates that FBP could not only report the flux in steady state, but also dynamically (Visser *et al.*, 2004; Bosch *et al.*, 2008).

In contrast, for example, the levels of ATP, ADP and AMP (obtained from van Meyenburg, 1969; Larsson *et al.*, 1997; Canelas *et al.*, 2011; Christen & Sauer, 2011), do not show any clear trend making it unlikely that the concentration of these metabolites would be the input for the respective regulatory machinery that controls the activity of the fermentative and respiratory pathways. This conclusion is further supported by findings from a recent study, in which mitochondrial NAD<sup>+</sup> carriers were deleted or overexpressed in *S. cerevisiae*. While these perturbations led to altered NAD and ATP levels, all the mutants switched from a fully respiratory metabolism to the respirofermentative one at the same glucose flux as the wild type (Vemuri *et al.*, 2007; Agrimi *et al.*, 2011), corroborating the idea that the levels of the energy cofactors levels are not likely to serve as flux signals.

How could the flux-information – imprinted into a metabolite's concentration (FBP, for example) – be coupled to the regulatory machinery to finally result in flux-dependent regulation? The first option for such coupling is an interaction of the flux-signaling metabolite directly with enzymes to activate or inactivate their activity (Figure 4), for which a couple of different mechanisms exist (Zorn & Wells, 2010). FBP, for example, is known to (i) activate pyruvate kinase (Otto *et al.*, 1986; Susan-Resiga & Nowak, 2003), (ii) have effects on the fructose 2,6-bisphosphate and AMP mediated activation of phosphofructokinase activity (Przybylski *et al.*, 1985) and (iii) inhibit oxidative phosphorylation through strongly inhibiting Complex IV (cytochrome c oxidase) and Complex III (ubiquinol:cytochrome c oxidoreductase) (Diaz-Ruiz *et al.*, 2008).

Alternatively, flux-signaling metabolites can be coupled to the regulatory machinery via interaction with signaling and regulatory proteins, such as kinases or transcription factors. Unlike in *E. coli*, for

which about a 100 metabolite-transcription factor interactions are known (ECOCYC database, Keseler *et al.*, 2011), we only know about a handful of such interactions in yeast, e.g. Gal3p, Put3p and Bas1p (for reviews see Sellick & Reece, 2005; Reece *et al.*, 2006; Campbell *et al.*, 2008). Here, the question is whether we simply do not know more (because they are hard to identify by classical biochemical means and as of today still no high-throughput method exists (Heinemann & Sauer, 2011) or whether they are not as prominent in yeast as they are in bacteria. Although most yeast transcription factors do not have small molecule binding pockets as, for example, the *E. coli* transcription factors (personal communication Nick Luscombe), novel metabolite-transcription factor interactions (Pinson *et al.*, 2009) and signaling protein interactions (Li & Snyder, 2011) were identified recently. Thus, there might be more to be discovered in the future. Similarly, we are also only about to realize that metabolite-binding RNA domains are also present in the genes of eukaryotes (Wachter, 2010), which might offer an alternative possibility how flux-signals imprinted into metabolite levels could be coupled to the regulatory machinery of a cell.

In contrast to sensing metabolic flux via flux-signaling metabolites, one could envision that certain enzymes directly sense metabolic flux. It is often speculated that hexokinase PII (Hxk2p) could do this, for example, via conformational changes that accompany the catalysis and that induce localization of Hxk2p to the nucleus or changes in a signaling complex of which Hxk2p is a component (Bisson & Kunathigan, 2003). All of this is however still elusive and as several sugars are not processed via hexokinase (such as galactose), a flux-sensor involving Hxk2 could not represent the full story.

### **III. Towards identifying the regulatory system that controls metabolism in flux-dependent manner**

In response to a sensed glycolytic flux, a cell must be able to diverge the flux into either the respiratory or fermentative pathways. Due to the regulatory system's global and likely complex architecture - overarching multiple cellular levels - we have currently only a few fragmented pieces of evidence for what the actual regulatory system could be:

1. The system needs to be fast. Glucose pulses in glucose- and ethanol-limited chemostat cultures result in an immediate onset of ethanol excretion (Visser *et al.*, 2004; Bosch *et al.*, 2008), which indicates that at least part of the system must reside on the fast enzymatic level, e.g. in the form of metabolite-enzyme interactions. One such action could be the recently identified inhibition of the respiratory chain through FBP (Diaz-Ruiz *et al.*, 2008). Alternatively, fast metabolite-kinase interactions could also be involved in modulation of enzyme activity through phosphorylation. Here, several new enzyme phosphorylation sites were recently identified (Breitkreutz *et al.*, 2010). With regard to the fact that the system needs to be fast, it is interesting to note that older findings with *in vitro* systems constituted of a small subset of glycolytic enzymes in a continuous system were found to show different states depending on the glucose feed rate (Schellenberger *et al.*, 1980).

2. Redox metabolism has likely a limited role in the system. Perturbations in the NAD<sup>+</sup> metabolism (which lead to changes in cellular NAD and ATP levels and affect growth rate) did not break the correlation between the sugar uptake and ethanol excretion rate (Agrimi *et al.*, 2011). Also increasing respiration or non-respiratory NADH oxidation in *S. cerevisiae* had only minor effects on the correlation between glycolytic flux and ethanol production rate (Vemuri *et al.*, 2007) (the data from these two references is displayed in Figure 1A, cf. supplementary information for details). Thus, although redox metabolism might be involved to a small extend in establishing overflow metabolism (Vemuri *et al.*, 2007), it does not seem to have a major influence on the distribution between respiratory and fermentative metabolism.

3. Genetic and chemical perturbations that block either the respiratory or the fermentative pathway result in complete system failures only at certain sugar uptake rates. For example, impinging on the capacity of the fermentative pathway by deletion of *ADHI* leads to a sick phenotype in glucose batch conditions (i.e. a high glucose uptake rate condition demanding fermentative metabolism), while this deletion has basically no phenotype on galactose (i.e. a low sugar uptake rate condition that demands a respiratory metabolism) (Ewald, Matt and Zamboni, 2011, unpublished). Vice versa, a deletion in the TCA cycle isoenzyme *SDHI*, which likely puts an upper limit on the flux through the TCA cycle, shows a practically lethal phenotype on galactose, while on glucose this mutation does not have a

phenotype (Ewald, Matt and Zamboni, 2011, unpublished). Along the same lines, blocking the respiratory chain with antimycin A only blocks growth in strains under conditions with a low sugar uptake rate (requiring a respiratory metabolism), while fermenting cells are not susceptible (Goffrini *et al.*, 2002; Fukuhara, 2003; Merico *et al.*, 2007). These observations suggest that when certain pathway cannot be used, because they are either chemically or genetically blocked, the regulatory system is not flexible enough to simply switch to the other metabolic mode.

## Conclusion

Currently, we have only a very limited understanding about the sensing and regulation system that - in *S. cerevisiae* and also in other yeast species - is responsible for the distribution of flux into the respiratory or fermentative pathways. What we basically know today is the following: (i) The system seems to regulate the metabolic phenotype in a glycolytic flux-dependent manner, which is basically independent of yeast species, growth conditions, sugar source and many different genetic and environmental perturbations and thus seems to be very robust. (cf. section I). (ii) The concentration of FBP (and likely also other metabolites) is flux-dependent making them ideally suited as potential flux-sensors (cf. section II). (iii) There are only very fragmented insights into the regulatory system's characteristics (cf. section III).

To eventually obtain a complete systems-level understanding of how eventually 'metabolic flux controls metabolic flux', it is clear that next to the classical tools of biological research also various systems biology approaches (Heinemann & Sauer, 2010) will be required. For example, high-throughput analytical technologies such as metabolomics and fluxomics will be required for identification of further potentially existing flux-signaling metabolites, analytical procedures for detection of novel small molecule-protein interactions and phospho-proteomics to further investigate the relevance of enzyme phosphorylation. Next to these discovery-driven applications of these modern -omics technologies, it will also be required to generate molecule abundance data which after subjecting them to computational top-down analyses will likely be able to extract regulatory interactions. Finally, bottom-up modeling approaches will be required to test hypothesis on whether

certain behavior can emerge from the quantitative and dynamic interaction of a select set of molecular players. Here, the grand challenge will be to find the right level of abstraction for the model overarching multiple levels of the cell. Only with a model available, it will ultimately be possible to show whether, and if yes how metabolic flux is sensed and used for regulation.

## **Conclusion**

To ultimately understand how the intricate system of the flux-sensing mechanism(s) and the respective regulatory machinery works, we will need to draw on all different branches of systems biology ranging from large-scale experimental approaches via top-down analyses to detailed modeling efforts.

## **Acknowledgements**

Funding is acknowledged from the SystemsX.ch project YeastX and from the Centre for Synthetic Biology of the University of Groningen. We would like to thank Nicola Zamboni and Jennifer Ewald for sharing data before publication and Guillermo Zampar for helpful comments on the manuscript.

## **References**

- Abbott DA, Knijnenburg TA, de Poorter LM, Reinders MJ, Pronk JT & van Maris AJ (2007) Generic and specific transcriptional responses to different weak organic acids in anaerobic chemostat cultures of *Saccharomyces cerevisiae* *FEMS Yeast Res* **7**: 819-833.
- Agrimi G, Brambilla L, Frascotti G, Pisano I, Porro D, Vai M & Palmieri L (2011) Deletion or overexpression of mitochondrial NAD<sup>+</sup> carriers in *Saccharomyces cerevisiae* alters cellular NAD and ATP contents and affects mitochondrial metabolism and the rate of glycolysis *Appl Environ Microbiol* **77**: 2239-2246.
- Aguilera J, Petit T, de Winde JH & Pronk JT (2005) Physiological and genome-wide transcriptional responses of *Saccharomyces cerevisiae* to high carbon dioxide concentrations *FEMS Yeast Res* **5**: 579-593.

363 Badotti F, Dario MG, Alves SL,Jr, Cordioli ML, Milette LC, de Araujo PS & Stambuk BU (2008)  
 364 Switching the mode of sucrose utilization by *Saccharomyces cerevisiae* *Microb Cell Fact* **7**: 4.

365 Barford JP & Hall RJ (1979) An examination of the Crabtree effect in *Saccharomyces cerevisiae*: the  
 366 role of respiratory adaptation *Microbiology* **114**: 267-275.

367 Basso TO, Dario MG, Tonso A, Stambuk BU & Gombert AK (2010) Insufficient uracil supply in fully  
 368 aerobic chemostat cultures of *Saccharomyces cerevisiae* leads to respiro-fermentative metabolism and  
 369 double nutrient-limitation *Biotechnol Lett* **32**: 973-977.

370 Beck C & von Meyenburg HK (1968) Enzyme pattern and aerobic growth of *Saccharomyces*  
 371 *cerevisiae* under various degrees of glucose limitation *J Bacteriol* **96**: 479-486.

372 Bisson LF & Kunathigan V (2003) On the trail of an elusive flux sensor *Res Microbiol* **154**: 603-610.

373 Blank LM, Kuepfer L & Sauer U (2005) Large-scale <sup>13</sup>C-flux analysis reveals mechanistic principles  
 374 of metabolic network robustness to null mutations in yeast *Genome Biol* **6**: R49.

375 Blom J, De Mattos MJ & Grivell LA (2000) Redirection of the respiro-fermentative flux distribution  
 376 in *Saccharomyces cerevisiae* by overexpression of the transcription factor Hap4p *Appl Environ*  
 377 *Microbiol* **66**: 1970-1973.

378 Boer VM, Crutchfield CA, Bradley PH, Botstein D & Rabinowitz JD (2010) Growth-limiting  
 379 intracellular metabolites in yeast growing under diverse nutrient limitations *Mol Biol Cell* **21**: 198-  
 380 211.

381 Boles E & Hollenberg CP (1997) The molecular genetics of hexose transport in yeasts *FEMS*  
 382 *Microbiol Rev* **21**: 85-111.

383 Bosch D, Johansson M, Ferndahl C, Franzen CJ, Larsson C & Gustafsson L (2008) Characterization  
 384 of glucose transport mutants of *Saccharomyces cerevisiae* during a nutritional upshift reveals a  
 385 correlation between metabolite levels and glycolytic flux *FEMS Yeast Res* **8**: 10-25.



386 Breitkreutz A, Choi H, Sharom JR *et al.* (2010) A global protein kinase and phosphatase interaction  
387 network in yeast *Science* **328**: 1043-1046.

388 Bull AT (2010) The renaissance of continuous culture in the post-genomics age *J Ind Microbiol*  
389 *Biotechnol* **37**: 993-1021.

390 Campbell RN, Leverentz MK, Ryan LA & Reece RJ (2008) Metabolic control of transcription:  
391 paradigms and lessons from *Saccharomyces cerevisiae* *Biochem J* **414**: 177-187.

392 Canelas AB, Ras C, ten Pierick A, van Gulik WM & Heijnen JJ (2011) An in vivo data-driven  
393 framework for classification and quantification of enzyme kinetics and determination of apparent  
394 thermodynamic data *Metab Eng* **13**: 294-306.

395 Christen S & Sauer U (2011) Intracellular characterization of aerobic glucose metabolism in seven  
396 yeast species by <sup>13</sup>C flux analysis and metabolomics *FEMS Yeast Res* **11**: 263-272.

397 Cordier H, Mendes F, Vasconcelos I & Francois JM (2007) A metabolic and genomic study of  
398 engineered *Saccharomyces cerevisiae* strains for high glycerol production *Metab Eng* **9**: 364-378.

399 Cortassa S & Aon M, A. (1998) The onset of fermentative metabolism in continuous cultures  
400 depends on the catabolite repression properties of *Saccharomyces cerevisiae* *Enzyme and Microbial*  
401 *Technology* **22**: 705-712.

402 Costenoble R, Picotti P, Reiter L, Stallmach R, Heinemann M, Sauer U & Aebersold R (2011)  
403 Comprehensive quantitative analysis of central carbon and amino-acid metabolism in *Saccharomyces*  
404 *cerevisiae* under multiple conditions by targeted proteomics *Mol Syst Biol* **7**: 464.

405 Daran-Lapujade P, Jansen ML, Daran JM, van Gulik W, de Winde JH & Pronk JT (2004) Role of  
406 transcriptional regulation in controlling fluxes in central carbon metabolism of *Saccharomyces*  
407 *cerevisiae*. A chemostat culture study *J Biol Chem* **279**: 9125-9138.

408 Daran-Lapujade P, Rossell S, van Gulik WM *et al.* (2007) The fluxes through glycolytic enzymes in  
 409 *Saccharomyces cerevisiae* are predominantly regulated at posttranscriptional levels *Proc Natl Acad Sci*  
 410 *U S A* **104**: 15753-15758.

411 De Deken RH (1966) The Crabtree effect: a regulatory system in yeast *J Gen Microbiol* **44**: 149-156.

412 de Kok S, Yilmaz D, Suir E, Pronk JT, Daran JM & van Maris AJ (2011) Increasing free-energy  
 413 (ATP) conservation in maltose-grown *Saccharomyces cerevisiae* by expression of a heterologous  
 414 maltose phosphorylase *Metab Eng* **13**: 518-526.

415 Diaz-Ruiz R, Averet N, Araiza D, Pinson B, Uribe-Carvajal S, Devin A & Rigoulet M (2008)  
 416 Mitochondrial oxidative phosphorylation is regulated by fructose 1,6-bisphosphate. A possible role in  
 417 Crabtree effect induction? *J Biol Chem* **283**: 26948-26955.

418 Diderich JA, Schepper M, van Hoek P *et al.* (1999) Glucose uptake kinetics and transcription of HXT  
 419 genes in chemostat cultures of *Saccharomyces cerevisiae* *J Biol Chem* **274**: 15350-15359.

420 Dietvorst J, Londesborough J & Steensma HY (2005) Maltotriose utilization in lager yeast strains:  
 421 MTT1 encodes a maltotriose transporter *Yeast* **22**: 775-788.

422 Does AL & Bisson LF (1989) Comparison of glucose uptake kinetics in different yeasts *J Bacteriol*  
 423 **171**: 1303-1308.

424 Elbing K, Larsson C, Bill RM, Albers E, Snoep JL, Boles E, Hohmann S & Gustafsson L (2004) Role  
 425 of hexose transport in control of glycolytic flux in *Saccharomyces cerevisiae* *Appl Environ Microbiol*  
 426 **70**: 5323-5330.

427 Fendt SM & Sauer U (2010) Transcriptional regulation of respiration in yeast metabolizing differently  
 428 repressive carbon substrates *BMC Syst Biol* **4**: 12.

429 Fendt SM, Buescher JM, Rudroff F, Picotti P, Zamboni N & Sauer U (2010) Tradeoff between  
 430 enzyme and metabolite efficiency maintains metabolic homeostasis upon perturbations in enzyme  
 431 capacity *Mol Syst Biol* **6**: 356.

432 Frick O & Wittmann C (2005) Characterization of the metabolic shift between oxidative and  
 433 fermentative growth in *Saccharomyces cerevisiae* by comparative <sup>13</sup>C flux analysis *Microb Cell Fact*  
 434 **4**: 30.

435 Fukuhara H (2003) The Kluver effect revisited *FEMS Yeast Res* **3**: 327-331.

436 Fung E, Wong WW, Suen JK, Bulter T, Lee SG & Liao JC (2005) A synthetic gene-metabolic  
 437 oscillator *Nature* **435**: 118-122.

438 Gamo FJ, Lafuente MJ & Gancedo C (1994) The mutation DGT1-1 decreases glucose transport and  
 439 alleviates carbon catabolite repression in *Saccharomyces cerevisiae* *J Bacteriol* **176**: 7423-7429.

440 Goffrini P, Ferrero I & Donnini C (2002) Respiration-dependent utilization of sugars in yeasts: a  
 441 determinant role for sugar transporters *J Bacteriol* **184**: 427-432.

442 Gombert AK, Moreira dos Santos M, Christensen B & Nielsen J (2001) Network identification and  
 443 flux quantification in the central metabolism of *Saccharomyces cerevisiae* under different conditions  
 444 of glucose repression *J Bacteriol* **183**: 1441-1451.

445 Ha SJ, Galazka JM, Kim SR, Choi JH, Yang X, Seo JH, Glass NL, Cate JH & Jin YS (2011)  
 446 Engineered *Saccharomyces cerevisiae* capable of simultaneous cellobiose and xylose fermentation  
 447 *Proc Natl Acad Sci U S A* **108**: 504-509.

448 Heinemann M & Sauer U (2011) From good old biochemical analyses to high-throughput omics  
 449 measurements and back *Curr Opin Biotechnol* **22**: 1-2.

450 Heinemann M & Sauer U (2010) Systems biology of microbial metabolism *Curr Opin Microbiol* **13**:  
 451 337-343.

452    Henricsson C, de Jesus Ferreira MC, Hedfalk K, Elbing K, Larsson C, Bill RM, Norbeck J, Hohmann  
 453    S & Gustafsson L (2005) Engineering of a novel *Saccharomyces cerevisiae* wine strain with a  
 454    respiratory phenotype at high external glucose concentrations *Appl Environ Microbiol* **71**: 6185-6192.

455    Heyland J, Fu J & Blank LM (2009) Correlation between TCA cycle flux and glucose uptake rate  
 456    during respiro-fermentative growth of *Saccharomyces cerevisiae* *Microbiology* **155**: 3827-3837.

457    Holsbeeks I, Lagatie O, Van Nuland A, Van de Velde S & Thevelein JM (2004) The eukaryotic  
 458    plasma membrane as a nutrient-sensing device *Trends Biochem Sci* **29**: 556-564.

459    Johnston M & Kim JH (2005) Glucose as a hormone: receptor-mediated glucose sensing in the yeast  
 460    *Saccharomyces cerevisiae* *Biochem Soc Trans* **33**: 247-252.

461    Jouhten P, Rintala E, Huuskonen A, Tamminen A, Toivari M, Wiebe M, Ruohonen L, Penttilä M &  
 462    Maaheimo H (2008) Oxygen dependence of metabolic fluxes and energy generation of *Saccharomyces*  
 463    *cerevisiae* CEN.PK113-1A *BMC Syst Biol* **2**: 60.

464    Keseler IM, Collado-Vides J, Santos-Zavaleta A *et al.* (2011) EcoCyc: a comprehensive database of  
 465    *Escherichia coli* biology *Nucleic Acids Res* **39**: D583-90.

466    Kotte O, Zaugg JB & Heinemann M (2010) Bacterial adaptation through distributed sensing of  
 467    metabolic fluxes *Mol Syst Biol* **6**: 355.

468    Kummel A, Ewald JC, Fendt SM, Jol SJ, Picotti P, Aebersold R, Sauer U, Zamboni N & Heinemann  
 469    M (2010) Differential glucose repression in common yeast strains in response to HXK2 deletion  
 470    *FEMS Yeast Res* **10**: 322-332.

471    Larsson C, Nilsson A, Blomberg A & Gustafsson L (1997) Glycolytic flux is conditionally correlated  
 472    with ATP concentration in *Saccharomyces cerevisiae*: a chemostat study under carbon- or nitrogen-  
 473    limiting conditions *J Bacteriol* **179**: 7243-7250.

474 Li X & Snyder M (2011) Metabolites as global regulators: a new view of protein regulation:  
 475 systematic investigation of metabolite-protein interactions may help bridge the gap between genome-  
 476 wide association studies and small molecule screening studies *Bioessays* **33**: 485-489.

477 Mahadevan R & Schilling CH (2003) The effects of alternate optimal solutions in constraint-based  
 478 genome-scale metabolic models *Metab Eng* **5**: 264-276.

479 Meijer MM, Boonstra J, Verkleij AJ & Verrips CT (1998) Glucose repression in *Saccharomyces*  
 480 *cerevisiae* is related to the glucose concentration rather than the glucose flux *J Biol Chem* **273**: 24102-  
 481 24107.

482 Merico A, Sulo P, Piskur J & Compagno C (2007) Fermentative lifestyle in yeasts belonging to the  
 483 *Saccharomyces* complex *FEBS J* **274**: 976-989.

484 Nissen TL, Schulze U, Nielsen J & Villadsen J (1997) Flux distributions in anaerobic, glucose-limited  
 485 continuous cultures of *Saccharomyces cerevisiae* *Microbiology* **143** ( Pt 1): 203-218.

486 Ostergaard S, Olsson L, Johnston M & Nielsen J (2000) Increasing galactose consumption by  
 487 *Saccharomyces cerevisiae* through metabolic engineering of the GAL gene regulatory network *Nat*  
 488 *Biotechnol* **18**: 1283-1286.

489 Otterstedt K, Larsson C, Bill RM, Stahlberg A, Boles E, Hohmann S & Gustafsson L (2004)  
 490 Switching the mode of metabolism in the yeast *Saccharomyces cerevisiae* *EMBO Rep* **5**: 532-537.

491 Otto A, Przybylski F, Nissler K, Schellenberger W & Hofmann E (1986) Kinetic effects of fructose-  
 492 1,6-bisphosphate on yeast phosphofructokinase *Biomed Biochim Acta* **45**: 865-875.

493 Peter Smits H, Hauf J, Muller S, Hobley TJ, Zimmermann FK, Hahn-Hagerdal B, Nielsen J & Olsson  
 494 L (2000) Simultaneous overexpression of enzymes of the lower part of glycolysis can enhance the  
 495 fermentative capacity of *Saccharomyces cerevisiae* *Yeast* **16**: 1325-1334.

496 Pinson B, Vaur S, Sagot I, Coulpier F, Lemoine S & Daignan-Fornier B (2009) Metabolic  
 497 intermediates selectively stimulate transcription factor interaction and modulate phosphate and purine  
 498 pathways *Genes Dev* **23**: 1399-1407.

499 Postmus J, Canelas AB, Bouwman J, Bakker BM, van Gulik W, de Mattos MJ, Brul S & Smits GJ  
 500 (2008) Quantitative analysis of the high temperature-induced glycolytic flux increase in  
 501 *Saccharomyces cerevisiae* reveals dominant metabolic regulation *J Biol Chem* **283**: 23524-23532.

502 Przybylski F, Otto A, Nissler K, Schellenberger W & Hofmann E (1985) Effects of fructose 1,6-  
 503 bisphosphate on the activation of yeast phosphofructokinase by fructose 2,6-bisphosphate and AMP  
 504 *Biochim Biophys Acta* **831**: 350-352.

505 Raab AM, Hlavacek V, Bolotina N & Lang C (2011) Shifting the fermentative/oxidative balance in  
 506 *Saccharomyces cerevisiae* by transcriptional deregulation of Snf1 via overexpression of the upstream  
 507 activating kinase Sak1p *Appl Environ Microbiol* **77**: 1981-1989.

508 Reece RJ, Beynon L, Holden S, Hughes AD, Rebora K & Sellick CA (2006) Nutrient-regulated gene  
 509 expression in eukaryotes *Biochem Soc Symp* (**73**): 85-96.

510 Reifengerger E, Freidel K & Ciriacy M (1995) Identification of novel HXT genes in *Saccharomyces*  
 511 *cerevisiae* reveals the impact of individual hexose transporters on glycolytic flux *Mol Microbiol* **16**:  
 512 157-167.

513 Rodriguez C & Flores C (2000) Mutations in GAL2 or GAL4 alleviate catabolite repression produced  
 514 by galactose in *Saccharomyces cerevisiae* *Enzyme Microb Technol* **26**: 748-755.

515 Rozpedowska E, Hellborg L, Ishchuk OP, Orhan F, Galafassi S, Merico A, Woolfit M, Compagno C  
 516 & Piskur J (2011) Parallel evolution of the make-accumulate-consume strategy in *Saccharomyces* and  
 517 *Dekkera* yeasts *Nat Commun* **2**: 302.

518 Rubio-Teixeira M, Van Zeebroeck G, Voordeckers K & Thevelein JM (2010) *Saccharomyces*  
 519 *cerevisiae* plasma membrane nutrient sensors and their role in PKA signaling *FEMS Yeast Res* **10**:  
 520 134-149.

521 Schellenberger W, Eschrich K & Hofmann E (1980) Selforganization of a glycolytic reconstituted  
 522 enzyme system: alternate stable stationary states, hysteretic transitions and stabilization of the energy  
 523 charge *Adv Enzyme Regul* **19**: 257-284.

524 Sellick CA & Reece RJ (2005) Eukaryotic transcription factors as direct nutrient sensors *Trends*  
 525 *Biochem Sci* **30**: 405-412.

526 Sierkstra LN, Nouwen NP, Verbakel JM & Verrips CT (1993) Regulation of glycolytic enzymes and  
 527 the Crabtree effect in galactose-limited continuous cultures of *Saccharomyces cerevisiae* *Yeast* **9**: 787-  
 528 795.

529 Sonnleitner B & Kappeli O (1986) Growth of *Saccharomyces cerevisiae* is controlled by its limited  
 530 respiratory capacity: Formulation and verification of a hypothesis *Biotechnol Bioeng* **28**: 927-937.

531 Susan-Resiga D & Nowak T (2003) The proton transfer step catalyzed by yeast pyruvate kinase *J Biol*  
 532 *Chem* **278**: 12660-12671.

533 Tai SL, Daran-Lapujade P, Luttik MA, Walsh MC, Diderich JA, Krijger GC, van Gulik WM, Pronk  
 534 JT & Daran JM (2007) Control of the glycolytic flux in *Saccharomyces cerevisiae* grown at low  
 535 temperature: a multi-level analysis in anaerobic chemostat cultures *J Biol Chem* **282**: 10243-10251.

536 van Hoek P, Flikweert MT, van der Aart QJ, Steensma HY, van Dijken JP & Pronk JT (1998) Effects  
 537 of pyruvate decarboxylase overproduction on flux distribution at the pyruvate branch point in  
 538 *Saccharomyces cerevisiae* *Appl Environ Microbiol* **64**: 2133-2140.

539 van Meyenburg K (1969) Katabolit-Repression und der Sprossungszyklus von *Saccharomyces*  
 540 *cerevisiae*. ETH Zurich, Zurich.

541 Van Urk H, Voll WS, Scheffers WA & Van Dijken JP (1990) Transient-state analysis of metabolic  
 542 fluxes in crabtree-positive and crabtree-negative yeasts *Appl Environ Microbiol* **56**: 281-287.

543 van Urk H, Postma E, Scheffers WA & van Dijken JP (1989) Glucose transport in crabtree-positive  
 544 and crabtree-negative yeasts *J Gen Microbiol* **135**: 2399-2406.

545 Velagapudi VR, Wittmann C, Schneider K & Heinzle E (2007) Metabolic flux screening of  
 546 *Saccharomyces cerevisiae* single knockout strains on glucose and galactose supports elucidation of  
 547 gene function *J Biotechnol* **132**: 395-404.

548 Vemuri GN, Eiteman MA, McEwen JE, Olsson L & Nielsen J (2007) Increasing NADH oxidation  
 549 reduces overflow metabolism in *Saccharomyces cerevisiae* *Proc Natl Acad Sci U S A* **104**: 2402-2407.

550 Verduyn C, Postma E, Scheffers WA & Van Dijken JP (1992) Effect of benzoic acid on metabolic  
 551 fluxes in yeasts: a continuous-culture study on the regulation of respiration and alcoholic fermentation  
 552 *Yeast* **8**: 501-517.

553 Verduyn C, Stouthamer AH, Scheffers WA & van Dijken JP (1991) A theoretical evaluation of  
 554 growth yields of yeasts *Antonie Van Leeuwenhoek* **59**: 49-63.

555 Verduyn C, Postma E, Scheffers WA & van Dijken JP (1990) Physiology of *Saccharomyces*  
 556 *cerevisiae* in anaerobic glucose-limited chemostat cultures *J Gen Microbiol* **136**: 395-403.

557 Visser D, van Zuylen GA, van Dam JC, Eman MR, Proll A, Ras C, Wu L, van Gulik WM & Heijnen  
 558 JJ (2004) Analysis of in vivo kinetics of glycolysis in aerobic *Saccharomyces cerevisiae* by application  
 559 of glucose and ethanol pulses *Biotechnol Bioeng* **88**: 157-167.

560 Wachter A (2010) Riboswitch-mediated control of gene expression in eukaryotes *RNA Biol* **7**: 67-76.

561 Weusthuis RA, Pronk JT, van den Broek PJ & van Dijken JP (1994) Chemostat cultivation as a tool  
 562 for studies on sugar transport in yeasts *Microbiol Rev* **58**: 616-630.



563 Wiebe MG, Rintala E, Tamminen A *et al.* (2008) Central carbon metabolism of *Saccharomyces*  
 564 *cerevisiae* in anaerobic, oxygen-limited and fully aerobic steady-state conditions and following a shift  
 565 to anaerobic conditions *FEMS Yeast Res* **8**: 140-154.

566 Wisselink HW, Toirkens MJ, del Rosario Franco Berriel M, Winkler AA, van Dijken JP, Pronk JT &  
 567 van Maris AJ (2007) Engineering of *Saccharomyces cerevisiae* for efficient anaerobic alcoholic  
 568 fermentation of L-arabinose *Appl Environ Microbiol* **73**: 4881-4891.

569 Ye L, Kruckeberg AL, Berden JA & van Dam K (1999) Growth and glucose repression are controlled  
 570 by glucose transport in *Saccharomyces cerevisiae* cells containing only one glucose transporter *J*  
 571 *Bacteriol* **181**: 4673-4675.

572 Youk H & van Oudenaarden A (2009) Growth landscape formed by perception and import of glucose  
 573 in yeast *Nature* **462**: 875-879.

574 Zastrow CR, Hollatz C, de Araujo PS & Stambuk BU (2001) Maltotriose fermentation by  
 575 *Saccharomyces cerevisiae* *J Ind Microbiol Biotechnol* **27**: 34-38.

576 Zhuang K, Vemuri GN & Mahadevan R (2011) Economics of membrane occupancy and respiration  
 577 fermentation *Mol Syst Biol* **7**: 500.

578 Zorn JA & Wells JA (2010) Turning enzymes ON with small molecules *Nat Chem Biol* **6**: 179-188.

579  
 580

581

## 582 **Figure legends**

583 **Figure 1:** (A) Ethanol production rate as a function of sugar uptake rate (normalized to c-mol); **inset:**  
584 wild type *S. cerevisiae* on glucose batch cultures (black open squares); wildtype *S. cerevisiae* on  
585 glucose limited chemostat cultures (black open triangles); *S. cerevisiae* hexose transporter mutant  
586 strains on glucose batch cultures (black open diamonds), arrows indicate measurement values of the  
587 TM6\* strain (see main text); *S. cerevisiae* glucose sensing mutant strains on glucose chemostat  
588 cultures (red circles); **main:** *S. cerevisiae* data from inset pooled (black squares); environmentally and  
589 genetically perturbed *S. cerevisiae* on glucose, incl. anaerobic conditions (blue triangles); *S. cerevisiae*  
590 wildtype and mutants on other sugars with different environmental conditions (red circles); other  
591 wildtype yeast species on glucose (green diamonds). (B) Ethanol production rate as a function of the  
592 growth rate; *S. cerevisiae* data from inset pooled (black squares); environmentally and genetically  
593 perturbed *S. cerevisiae* on glucose, incl. anaerobic conditions (blue triangles); *S. cerevisiae* wildtype  
594 and mutants on other sugars with different environmental conditions (red circles); other wildtype yeast  
595 species on glucose (green diamonds). In some cases, data was not available for both, the ethanol  
596 production rate and the growth rate. The references for the respective data points are provided in the  
597 supplement.

598 **Figure 2:** (A) The collected data in Fig. 1A suggest that there is a system (indicated by the black box)  
599 in place that depending on the sugar uptake rate generates different metabolic phenotypes in terms of  
600 fermentative and respiratory activity. (B) This system likely comprises of a flux-sensor that connects a  
601 rate  $r$  to a state  $x$ , which is received by a controller that in turn realizes the necessary regulatory  
602 adjustments in a flux-dependent manner.

603 **Figure 3:** Glucose isomerase flux as a function of the sugar uptake rate (normalized to c-mol); data  
604 from *S. cerevisiae* on glucose batch and chemostat cultures (black squares); environmentally and  
605 genetically perturbed *S. cerevisiae* on glucose (blue triangles); *S. cerevisiae* on galactose batch  
606 conditions (red circles); other yeast species on glucose batch conditions (green diamonds).

**Figure 4:** Metabolite levels as a function of sugar uptake rate (normalized to c-mols). **(A)** Fructose-1,6-bisphosphate concentrations levels, **(B)** ATP concentration levels, **(C)** ADP concentration levels, **(D)** AMP concentration levels. *S. cerevisiae* glucose batch and chemostat cultures (black squares), *S. cerevisiae* environmental perturbation (blue triangles), other yeast species on glucose batch cultures (green diamonds).

**Figure 5:** Different manners in which flux-information imprinted into concentration levels of flux-signaling metabolites could be coupled to the regulatory machinery.